

Addendum to “Coherent radio pulses from GEANT generated electromagnetic showers in ice”

Soebur Razzaque,¹ Surujhdeo Seunarine,² Scott W. Chambers,³ David Z. Besson,³ Douglas W. McKay,³ John P. Ralston,³ and David Seckel⁴

¹*Department of Astronomy & Astrophysics, Penn State University, University Park, Pennsylvania 16802*

²*Department of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand*

³*Department of Physics & Astronomy, University of Kansas, Lawrence, Kansas 66045 and*

⁴*Bartol Research Institute, University of Delaware, Newark, Delaware 19716*

(Dated: February 2, 2008)

We reevaluate our published calculations of electromagnetic showers generated by GEANT 3.21 and the radio frequency pulses they produce in ice. We are prompted by a recent report showing that GEANT 3.21-modeled showers are sensitive to internal settings in the electron tracking subroutine. We report the shower and pulse characteristics obtained with different settings of GEANT 3.21 and with GEANT 4. The default setting of electron tracking in GEANT 3.21 we used in previous work speeds up the shower simulation at the cost of information near the end of the tracks. We find that settings tracking e^- and e^+ to lower energy yield a more accurate calculation, a more intense shower, and proportionately stronger radio pulses at low frequencies. At high frequencies the relation between shower tracking algorithm and pulse spectrum is more complex. We obtain radial distributions of shower particles and phase distributions of pulses from 100 GeV showers that are consistent with our published results.

PACS numbers: 96.40.Pq, 95.85.Bh, 95.85.Ry, 29.40.-n

In a recent paper [1] we reported the results of an extensive study of the radio frequency pulse emitted by an electron-induced electromagnetic shower. Application to radio signal detection of ultrahigh energy electron neutrino interaction in ice provided our motivation [2]. We used the standard particle physics detector-simulation package GEANT 3.21 (G3 afterwards) [3] with default settings to generate the showers. Our total track-lengths of electrons and number of electrons at shower maximum were 68% of those in the paper by Zas, Halzen and Stanev (ZHS) [4]. Our pulse height at the Cherenkov angle at 1 GHz was 72% of the ZHS result. Our study of sensitivity of shower development to reasonable changes in input cross sections showed no effects large enough to explain the difference between G3 and ZHS shower profiles and track lengths.

The source of the bulk of the discrepancy was recently presented by Alvarez-Müniz, Marquéz, Vázquez and Zas [5]. They noticed that the default setting of a parameter called “ABAN” in the electron tracking subroutine of G3, the setting used in our studies, gives shorter total, projected and weighted track lengths and lower particle yield at shower maximum than GEANT 4 (G4 afterwards) or G3 with a choice of tracking rules different from the default. Their results for track-length and shower profile from the latter simulations are 90% of those of the original ZHS work for these quantities and in closer agreement than G3 showers with the default setting [1, 5].

Track lengths are one indicator of the strength of electric field pulses to be expected from the shower. However the electric field at a given angle and frequency depends on a combination of many factors. We make a comparison among the shower *and* electric field pulse results from G3 (with several different combinations of electron track-

ing rules) and G4. We report results for shower profiles, total track lengths, energy loss (dE/dx) and radial distribution of the number of electrons minus the number of positrons around the shower maximum. We tabulate the G3, G4 and ZHS track-lengths and yields at shower maximum for ease of comparison of shower results. Regarding the electric field in the Fraunhofer regime, we report radio pulse strength at the Cherenkov angle and the phase behavior of the radio pulse both on and off the Cherenkov peak. We find that the particle yield and charge imbalance are larger with improved tracking. At low frequencies the pulses emitted at the Cherenkov angle are correspondingly stronger as well.

Electron tracking in GEANT: The subroutine responsible for tracking electrons in G3 is called GTELEC. In its default setting, ABAN=1, if the electron range is too short to reach the boundary of a detector volume, and the distance to its next bremsstrahlung emission is greater than its range, the electron’s energy is deposited in the volume and it is stopped. This is done regardless of the value of the electron kinetic energy threshold parameter CUTELE set by a user in the input card file. This was presumably made the default to increase the calculation speed of high energy particle physics detector simulations in non-sensitive volumes, where detailed tracking was not needed [6]. With the default setting, however, track lengths are reduced significantly, with a consequent reduction in the calculated radio Cherenkov emission. This is what we observed in our previous study [1], where we compared results from G3 with default setting to the results of [4].

In practice, it seems for applications such as ours the preferred [7] setting is ABAN=0, which tracks all electrons down to the threshold set by the user. The setting

ABAN=2 allows the user to track electrons precisely (similar to ABAN=0 setting) in sensitive volumes or not so precisely in the non-sensitive volumes [8]. “Sensitive” vs. “non-sensitive” settings are distinguished by the value of a step parameter called “STMIN” in the tracking medium parameter list, where “sensitive” produces many more steps per track than “non-sensitive”. G4 has range cuts that correspond to energy thresholds for the production of particles. A particle, once produced, is tracked down to rest regardless of its production threshold.

Results: We expand on the study reported in Ref. [5], where showers produced with G3 ABAN=1, G3 ABAN=2 with “sensitive” volume, G4 and ZHS are compared. We run G3 for four different tracking choices, namely ABAN=0, ABAN=1, ABAN=2 sensitive volume (S) and ABAN=2 non-sensitive volume (NS) settings, to create showers in ice. We run G4 for comparison as well. We then repeat our study [1] of the key features of the showers and the consequent key features of the radio pulses: pulse angular distributions, phase distributions and frequency spectra.

1) *Shower Characteristics:* First we investigate the important issue of total track lengths in Table I. We note that the track lengths for ABAN=2 (NS) setting are similar to the default ABAN=1 setting; the ABAN=2 (S) setting used in Ref. [5] gives results similar to the preferred ABAN=0 setting and both are similar to the results from G4. The agreement between track lengths produced by these latter three GEANT versions and those produced by the ZHS code improves as one compares total, total projected $e + p$ and total projected $e - p$ track lengths. Comparing the projected $e + p$ and $e - p$ track lengths, we see that the loss of track length for ABAN=1 is more significant for electrons than for positrons. We expect this to be the case, since a large part of the electron excess comes from low energy electrons swept into the shower by Compton scattering and δ -ray production.

Despite the improved agreement on total track lengths, there are still significant differences between the G3 ABAN=0, G3 ABAN=2(S), and G4 simulations. For each code individually, however, the variance in total track length 36 – 40 cm is reasonable. For a given shower energy E_0 , the total track length L_{tot} is determined primarily by the rate of ionization energy losses, where the energy transfer is less than a low threshold $E_t \sim 100$ keV. Then $L_{tot} \approx E_0 / (dE/dx)_{ion}$, as expected under the conditions that a) soft ionization losses are essentially independent of particle energy and b) hard interactions do not “remove” energy from the shower. Hard δ -rays create their own track length and hard γ ’s return their energy to the shower in pair production and Compton scattering processes. Some energy is lost from the shower as soft particles fall below E_t . The fluctuations in track length are thus determined by the fluctuations in the number of soft particles multiplied by $E_t / (dE/dx)_{ion}$. Estimating the number of soft particles by $N_t = E_0 / E_t$, the fluctuations in total track length are of order $\delta L_{tot} \sim E_t / (dE/dx)_{ion} \sqrt{N_t} = \sqrt{E_0 E_t} / (dE/dx)_{ion} = 50$ cm, for

$E_0 = 100$ GeV, $E_t = 100$ keV, and $(dE/dx)_{ion} = 2$ MeV/cm. Although this is a rough estimate, the observed values of σ for L_{tot} are consistent with expectations for the newer simulations.

This estimate leads to several observations. a) The deviations in L_{tot} between G3 ABAN=0, G3 ABAN=2(S), and G4 are small but significant; this is under study. b) Due to the default decision in the tracking code to abandon some particles, the older G3 simulations have an effective value of $E_t \sim 5$ MeV, which implies $\delta L \sim 3.5$ m, generally consistent with the Table entries. c) For $e - p$ track length, the fluctuations are dominated by the competition between pair production and Compton scattering, for which one may estimate $E_t \sim 20$ MeV and $\delta L \sim 7$ m.

Particle yields at the maxima of longitudinal shower profiles are also listed in Table I. The results from the ZHS code, G3 with ABAN=0, ABAN=2 with sensitive volume, and G4 all agree reasonably well. Agreement is especially good for the excess charge, where results agree within errors. Again, the low yields from G3 ABAN=1 and ABAN=2 (NS) are consistent with the interpretation that lost track length is caused by a deficiency of low energy electrons being scattered into the shower.

Next we consider how individual particle tracks contribute to the radio pulse. For low frequencies it can be shown that the contributions are in phase and the pulse is proportional to the weighted, projected track length, labeled (e-p) in Table 1. As the frequency increases, this simple coherence breaks down as described by two parameters, which we called [1] the Cherenkov Phase (CP) and the Translational Phase (TP). The CP is related to the phase coherence within a single track segment, whereas the TP describes the relative phase among track segments due to their different positions within the shower, relative to the observer.

In general, we find that the different simulations produce phase distributions similar to those shown in Fig. 17 of Ref. [1]. Specifically, the radial charge distribution at shower maximum, shown to peak at $\rho \simeq 0.5$ cm in Fig. 22 of Ref. [1], is reproduced apart from an overall increase in normalization [9]. Moreover, the radial charge distribution early in the shower evolution has a core dominated by high energy particles, and the normalization of this core does not depend on the details of the electron tracking at low energy.

2) *Radio frequency pulse:* Next we examine how the changes in shower simulation affect the modeled radio Cherenkov pulses. At low frequencies, all particles contribute in phase, so the increase in the excess track length found in G3 ABAN=0 and G4 should be reflected by an increase in the amplitudes of the low frequency part of the spectrum as compared to showers produced with G3 ABAN=1. In Fig. 1 we see that this is indeed the case, where the spectrum $R \times |\vec{E}|$ is shown for an observer on the Cherenkov cone.

In Fig.2 we extend the frequency scale out to 10 GHz, which suffices to show clearly a transition from low to

TABLE I: Track length and particle yield results from an average of 100 GeV electron induced shower using different Monte Carlo shower codes. The error bars correspond to error in the mean, $s/\sqrt{N}$, where s is the standard deviation reported within parentheses and N is the number of showers used in each case.

Monte Carlo shower code	No. of showers	Total track lengths			Particle yield at shower max		
		Absolute ($e + p$) [m]	Projected ($e + p$) [m]	Projected ($e - p$) [m]	Total ($e + p$)	Excess ($e - p$)	Excess fraction
G3 (ABAN=0) ^a	20	542.74 $\pm$ 0.08 (0.36)	455.3 $\pm$ 0.2 (0.9)	125.0 $\pm$ 2.0 (9.0)	148 $\pm$ 5 (22)	42 $\pm$ 3 (13)	28%
G3 (ABAN=1) ^b	20	389.51 $\pm$ 0.48 (2.16)	365.7 $\pm$ 0.5 (2.2)	76.3 $\pm$ 1.5 (6.7)	111 $\pm$ 7 (31)	20 $\pm$ 2 (9)	18%
G3 (ABAN=2, NS)	20	389.40 $\pm$ 0.71 (3.18)	367.0 $\pm$ 0.7 (3.1)	78.4 $\pm$ 2.8 (12.5)	113 $\pm$ 10 (45)	25 $\pm$ 5 (22)	22%
G3 (ABAN=2, S) ^c	20	561.05 $\pm$ 0.09 (0.40)	459.8 $\pm$ 0.3 (1.3)	125.8 $\pm$ 1.9 (8.5)	148 $\pm$ 8 (36)	41 $\pm$ 3 (13)	28%
G4	100	572.58 $\pm$ 0.04 (0.40)	466.3 $\pm$ 0.2 (1.6)	135.0 $\pm$ 0.8 (7.9)	153 $\pm$ 3 (27)	45 $\pm$ 1 (13)	30%
ZHS	20	642.17 $\pm$ 0.06 (0.25)	516.6 $\pm$ 0.2 (0.9)	135.2 $\pm$ 1.5 (6.8)	164 $\pm$ 6 (28)	44 $\pm$ 2 (11)	27%

^aPreferred setting (new).

^bDefault setting (old).

^cSetting used in Ref. [5] (similar to ABAN=0).

high frequency for the same sets of simulations. In this range, the spectra for the G3 ABAN=0 and G4 showers flatten out, but the G3 ABAN=1 spectrum grows slowly, so that the difference in their normalizations, relative to G4, is less than 20% for $f = 10$ GHz. We infer that in this region the spectra are not simply described by the phase coherent sum of amplitudes from a large number of low energy electrons. We note two effects that could be important in this region. First, a relatively small number of high energy particles near the shower axis could contribute strongly to the spectrum in this region. Since this part of the shower evolution is not sensitive to low energy electron tracking, we may see the G3 and G4 results converging. This trend continues at higher frequency. Second, a large fraction of the track-segment amplitudes are no longer coherent. For that fraction, the magnitude of $\vec{E}$ increases as a random walk as the track length increases. In this case, one would expect the amplitude to increase as the square root of the number of track segments, rather than linearly, which reduces the ratio of the pulse from the G4 generated shower to that from ABAN=1 as the frequency increases. This is seen clearly in Figs. 1 and 2. Given that total and projected track lengths are proportional to energy [1, 4], the effects just outlined should be relevant when comparing pulses from showers of different energies using the same simulation.

Figure 3 shows a comparison between pulse spectra from 1 TeV showers and 100 GeV showers, both generated by G4. The spectra are extended up to 50 GHz. For frequencies above 15 GHz, the results are fully consistent with Fig. 25 of Ref. [1]. At low frequencies, below a few GHz, the results are consistent as well, after accounting for the change in normalization between G4 and G3 ABAN=1. At 50 GHz, the TeV average, scaled down by a factor of 10, is smaller by roughly a factor $3 \sim \sqrt{10}$ than the 100 GeV average, possibly hinting that the “random walk” increase of $\vec{E}$ with energy, and therefore total track length, is setting in. At the same time, it seems clear that a combination of effects must be in play for intermediate frequencies.




FIG. 1: Direct Monte Carlo calculation of the low frequency end of the spectrum of the average electric field amplitude for twenty 100 GeV showers. The showers are run using G4 and G3 with both the preferred ABAN=0 and the default ABAN=1 settings. The same code was used to calculate electric fields from all the GEANT showers.

Since the flat spectra observed above several GHz contrast with the sharp falloff expected above the 1-2 GHz scale [4, 10], we referred to this phenomenon as an “extended coherence regime” in [1]. This is different from the use of coherence to refer to the rapid rise of the electric field strength as frequency increases in the region where frequency is low compared to the scale set by the peak at 0.5 cm in the shower radial distribution.

Summarizing, we see that the improved electron tracking performed by G4 or G3 ABAN=0 results in proportionately increased radio Cherenkov spectra at frequencies below a few GHz. At high frequencies the situation is more complicated. We are conducting further studies at a variety of energies and including showers initiated by γ -rays to identify the determining factors.

Summary and further discussion of results:

Motivated by the observation in [5] that the default setting for electron tracking in G3 stops many electrons prematurely, we have reviewed the calculations reported in [1]. In [1] we used G3 with default setting and found




FIG. 2: Direct Monte Carlo calculation of the frequency spectrum of the average electric field amplitude for twenty 100 GeV showers. The showers are run by G4 and G3 with both the preferred ABAN=0 and the default ABAN=1 settings. The same code was used to calculate electric fields for all the GEANT showers.




FIG. 3: Direct Monte Carlo calculation of the frequency spectrum of the average electric field magnitude for one hundred 100 GeV showers and ten 1 TeV showers run by G4. The 1 TeV spectrum is divided by a scaling factor of 10.

a persistently smaller shower and a weaker electromagnetic pulse at low frequencies than those reported in [4]. We find that this discrepancy largely disappears when electrons are tracked more completely in G3 or when us-

ing G4 generated showers, similar to results reported in [5]. Figure 1 shows that the strength of the pulses at low frequency clearly reflect the track lengths and particle yield at shower maximum. The high frequency behavior is more complex and is under further study.

The particles in the electromagnetic showers in a uniform medium, ice in the case of interest here, should be tracked as long as they still contribute to the Cherenkov signal. This is the case, for the ZHS, G4 and G3 simulations with the tracking option ABAN=0 or ABAN=2 with sensitive volume; the results from all these simulations agree at the 10% level. There is physics, namely significant emission, that is lost if the tracks are stopped prematurely and their energy dumped into the medium. Though choosing the default (ABAN=1) setting affects only the low energy tail of the shower, below the few MeV range, the population of particles (predominately electrons) is so high at the low energy end that the field contribution cannot be ignored.

For applications of the shower and pulse codes to neutrino detection experiments, the difference is obviously important. For example, below 1 GHz where currently deployed antennas are sensitive, the expected radio emission pulse is 1.3 to 1.4 times higher using the G3 or G4 code with particles tracked to threshold compared to the truncated showers produced by G3 with the ABAN=1 setting. The more correct treatments that follow the particles to lower energy produce significantly larger fields on the Cherenkov cone. This increases the effective volume estimates and leads to higher sensitivity estimates for cosmic neutrino detection systems like the Radio Ice Cherenkov Experiment (RICE) [11] and the recently approved balloon-borne experiment ANITA [12], both using the Antarctic polar ice cap as a neutrino target.

Acknowledgements: We thank J. Le Vaillant and J. Adams for help running GEANT simulations. We acknowledge helpful communications with M. Maire, J. Alvarez-Muñiz, E. Marqués and E. Zas. Work was supported in part by the NSF, the DOE and the University of Kansas General Research Fund. S. Seunarine was supported by a Marsden grant. S. Razzaque was partially supported by NSF AST0098416 grant.

-
- [1] S. Razzaque et al., Phys. Rev. D **65**, 103002 (2002).
 - [2] D. Besson in *RADHEP 2000*, Los Angeles, D. Salzberg and P. Gorham Eds., AIP Conference Proceedings, Vol. 579, pg. 157 (2001).
 - [3] R. Brun et al., GEANT 3 Manual, CERN Program Library Long Writeup W5013, 1994.
 - [4] E. Zas, F. Halzen and T. Stanev, Phys. Rev. D **45**, 362 (1992).
 - [5] J. Alvarez-Muñiz, E. Marqués, R. Vázquez and E. Zas, astro-ph/0206043.
 - [6] For historical reasons, the ABAN choices and their application are not described in the user's manual [3].
 - [7] We thank Michel Maire for communication about the

- electron tracking treatments in GEANT 3.21.
- [8] See parameters ISVOL, DEEMAX and STMIN under CONS200 section of the GEANT manual [3].
- [9] R. Buniy and J. Ralston, Phys. Rev. D **65**, 016003 (2002).
- [10] J. Alvarez-Muñiz, R. Vázquez and E. Zas, Phys. Rev. D **62**, 063001 (2000).
- [11] I. Kravchenko et al., Astroparticle Physics **19**, 15 (2003); I. Kravchenko et al., astro-ph/0206371 (Astroparticle Physics, in press).
- [12] P. Gorham, in *Ultra High Energy Particles from Space*, Aspen (2002).